

-[®]
(BRONCHO-MUNAL[®])

Indication: 1 3,5 *Haemophilus influenzae, Streptococcus (Diplococcus) pneumoniae, Klebsiella pneumoniae and ozaenae, Staphylococcus aureus, Streptococcus pyogenes and viridans, Moraxella (Branhamella/Neisseria) catarrhalis;*

Contraindications: (132), (171), (421), ;

Side effects: 3 - ;

Other: R07AX.

Interactions: *Streptococcus pneumoniae, Haemophilus influenzae, Klebsiella pneumoniae, Klebsiella ozaenae, Staphylococcus aureus, Streptococcus viridans, Streptococcus pyogenes, Moraxella catarrhalis.*

— CD4 - ;

— IgA ;

— IgE ;

— IgA ;

1 6 12
20 10

6

1 (23)/

4

6

,3

20

10

(, ,).

12

6

®

7

/100 < 1/10), (1/1000 < 1/100), (1/10000 < 1/1000), edDRA (< 1/10000), : (1/10), (1

5

25°

10

, 1 (10 × 1) 3 (10 × 3)

.

•

. ., / Lek Pharmaceuticals d. d., Slovenia ().

57, 1526, / Verovskova 57, 1526 Ljubljana, Slovenia.

•